

Name: \_\_\_\_\_

**Practice Exam 1**  
**CS 591Q/791V - Pattern Recognition**  
Total points: 50  
Posted on: March 18, 2009

Please read this first:

- ⓘ This is an in-class, closed-book exam consisting of 4 questions. ☺
- ⓘ In order for your exam to be graded you will have to turn in your solutions by 12:30pm.
- ⓘ You are not permitted to engage in *any* kind of discussion during the exam.
- ⓘ If a question seems ambiguous, please state your assumption and proceed to solve it.
- ⓘ An act of academic dishonesty will fetch you a 0 in the exam. ☹

- 
1. [15 points] Briefly explain the following terms: (a) Bootstrapping; (b) Parzen window; (c) Fisher's linear discriminant.
  2. [10 points] Consider a set of one-dimensional values sampled from an unknown density  $p(x)$ : 1, 1.5, 1.75, 2, 2.5, 2.75, 3, 5, 6, 6.25, 6.5, 7, 7.5. Estimate the value of the density function,  $\hat{p}(x)$ , at  $x = 0, 1, 3, 5, 7, 9$ , by using the parzen window technique with a *uniform* kernel function of window width 1.
  3. [10 points] Suppose we have three classes ( $C_1, C_2, C_3$ ) with  $P(C_1) = 0.5$ ,  $P(C_2) = 0.25$  and  $P(C_3) = 0.25$  and the following distributions

$$p(x|C_1) \sim N(0, 1); \quad p(x|C_2) \sim N(0.5, 1); \quad p(x|C_3) \sim N(1, 1)$$

and that we sample the following four points:  $x = 0.6, 0.1, 0.9, 1.1$ . Calculate explicitly the probability that the sequence actually came from  $C_1, C_3, C_3, C_2$ . Be careful to consider normalization.

4. [15 points] Let  $p(x|C_1) \sim N(\mu_1, \sigma_1^2)$  and  $p(x|C_2) \sim N(\mu_2, \sigma_2^2)$ , where the density parameters for both categories are unknown. Assume  $P(C_1) = P(C_2) = 0.5$ . Suppose the following training sets  $D_1$  and  $D_2$  are available for classes  $C_1$  and  $C_2$ , respectively:

$$D_1 = \{0.67, 1.19, -1.20, -0.02, -0.16\},$$

$$D_2 = \{1.00, 0.55, 2.55, -1.65, 1.61\}.$$

- (a) Find the maximum likelihood estimates for the four unknown parameters based on the training sets  $D_1$  and  $D_2$ .
  - (b) What is the Bayes decision rule using the estimated values for the parameters? According to the Bayes decision rule, to which class would you assign the test pattern  $x = 0.5$ ?
  - (c) What is the Bayes error rate using the estimated values for the parameters?
-